

Groups and Representations

Homework Assignment 7 (due on 6 December 2023)

Problem 25

We consider the abelian group $C_3 = \{e, a, a^{-1}\} \cong \mathbb{Z}_3$.

- a) How many (non-equivalent) irreps does C_3 have, what are their dimensions and how often do they appear in the regular rep?
- b) Show that

$$e_1 = \frac{1}{3}(e + a + a^{-1})$$

is a primitive idempotent, generating the trivial rep.

- c) Use the ansatz

$$e_2 = xe + ya + za^{-1}$$

in order to find all primitive idempotents.

- d) For each primitive idempotent find out whether it generates a new (non-equivalent) irrep or an irrep equivalent to one generated by a previous idempotent.
- e) Specify all minimal left ideals and construct the corresponding irreps of C_3 . Collect your results in a table.

Problem 26

Let V be a vector space and $A : V \rightarrow V$ a linear map. Show that if A is nilpotent (i.e. if for some $n \in \mathbb{N}$ we have $A^n v = 0 \forall v \in V$) then $\text{tr } A = 0$.

Problem 27

For $A \in \mathbb{C}^{n \times n}$ the matrix exponential is defined as

$$e^A = \exp(A) = \sum_{\nu=0}^{\infty} \frac{A^\nu}{\nu!}.$$

Prove:

- a) The series converges absolutely and uniformly.

HINT: On $\mathbb{C}^{n \times n}$ use the operator norm

$$\|A\| = \sup_{v \in \mathbb{C}^n \setminus \{0\}} \frac{|Av|}{|v|},$$

for which we have $\|AB\| \leq \|A\| \|B\|$.

- b) For $T \in GL(n)$ we have $e^{TAT^{-1}} = Te^AT^{-1}$.
 c) e^{tA} is the unique solution of the initial value problem $\dot{X}(t) = AX(t)$, $X(0) = \mathbb{1}$.
 d) For $t, s \in \mathbb{C}$ we have $e^{(t+s)A} = e^{tA}e^{sA}$.
 e) $(e^A)^\dagger = e^{(A^\dagger)}$.
 f) $\det e^A = e^{\text{tr} A}$.

Problem 28³

We can also write elements of the $\mathcal{A}(S_n)$ in birdtrack notation. In particular, we denote symmetrisers and anti-symmetrisers by open and solid bars, respectively, i.e.

$$\frac{1}{n!} s = \frac{1}{n!} \sum_{p \in S_n} p = \begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \\ \vdots \\ \text{---} \\ | \\ \text{---} \\ | \\ \vdots \\ \text{---} \end{array} \quad \text{and} \quad \frac{1}{n!} a = \frac{1}{n!} \sum_{p \in S_n} \text{sgn}(p)p = \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \\ \vdots \\ \text{---} \\ | \\ \text{---} \\ | \\ \vdots \\ \text{---} \end{array}.$$

Note that we include a factor of $\frac{1}{n!}$ in the definition of bars over n lines. For instance,

$$\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \frac{1}{2} (\text{---} + \text{---}) \quad \text{and} \quad \begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array} = \frac{1}{3!} (\text{---} - \text{---} - \text{---} - \text{---} + \text{---} + \text{---}). \tag{*}$$

Notice that in birdtrack notation the sign of a permutation, $(-1)^K$, is determined by the number K of line crossings; if more than two lines cross in a point, one should slightly perturb the diagram before counting, e.g. $\text{---} \rightsquigarrow \text{---}$ ($K=3$).

- a) Expand $\begin{array}{c} \text{---} \\ \text{---} \\ \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \end{array}$ and $\begin{array}{c} \text{---} \\ | \\ \text{---} \\ | \\ \text{---} \\ | \\ \vdots \\ \text{---} \\ | \\ \text{---} \\ | \\ \vdots \\ \text{---} \end{array}$ as in (*).

³will be discusses in the lecture on Thu 30 Nov 2023

We also use the corresponding notation for partial (anti-)symmetrisation over a subset of lines, e.g.

$$\begin{aligned} \text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} &= \frac{1}{2} \left(\text{---} \text{---} \text{---} + \text{---} \text{---} \text{---} \right) \quad \text{or} \\ \text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} &= \frac{1}{2} \left(\text{---} \text{---} \text{---} - \text{---} \text{---} \text{---} \right) = \frac{1}{2} \left(\text{---} \text{---} - \text{---} \text{---} \right). \end{aligned}$$

It follows directly from the definition of S and A that when intertwining any two lines S remains invariant and A changes by a factor of (-1) , i.e.

$$\text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} = \text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} \quad \text{and} \quad \text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} = - \text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---}.$$

- b) Explain why this implies that whenever two (or more) lines connect a symmetriser to an anti-symmetriser the whole expression vanishes, e.g.

$$\text{---} \begin{array}{|c} \hline \\ \hline \end{array} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} = 0.$$

Symmetrisers and anti-symmetrisers can be built recursively. To this end notice that on the r.h.s. of

$$\text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} = \frac{1}{n} \left(\text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} + \text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} + \dots + \text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} \right)$$

we have sorted the terms according to where the last line is mapped – to the n th, to the $(n-1)$ th, \dots , to the first line line. Multiplying with $\text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---}$ from the left and disentangling lines we obtain the compact relation

$$\text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} = \frac{1}{n} \left(\text{---} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} + (n-1) \text{---} \begin{array}{|c} \hline \\ \hline \end{array} \begin{array}{|c} \hline \\ \hline \end{array} \text{---} \right).$$

- c) Derive the corresponding recursion relation for anti-symmetrisers.