

FOUNDATIONS OF QUANTUM MECHANICS: ASSIGNMENT 10

Exercise 38: Essay question. (15 points)

Explain why Schrödinger's theory Sm has a many-worlds character.

Exercise 39: Marginal and conditional distribution (18 points)

Consider two random variables X, Y that assume only values ± 1 . Their joint distribution can be described by a 2×2 table of probabilities. **(a)** Give a generic example of such a table (i.e., one without symmetries). For your table, compute **(b)** the marginal distribution of X and **(c)** that of Y , as well as **(d)** the conditional distribution of X , given that $Y = +1$, **(e)** the expectation value $\mathbb{E}(X)$, and **(f)** $\mathbb{E}(XY)$.

Exercise 40: Spin singlet state (17 points)

Verify through direct computation that in the spin space $\mathbb{C}^4 = \mathbb{C}^2 \otimes \mathbb{C}^2$ of two spin- $\frac{1}{2}$ particles,

$$\begin{aligned} & |x\text{-up}\rangle|x\text{-down}\rangle - |x\text{-down}\rangle|x\text{-up}\rangle \\ &= |y\text{-up}\rangle|y\text{-down}\rangle - |y\text{-down}\rangle|y\text{-up}\rangle \\ &= |z\text{-up}\rangle|z\text{-down}\rangle - |z\text{-down}\rangle|z\text{-up}\rangle \end{aligned} \tag{1}$$

up to phase factors.

Exercise 41: Distinguish ensembles (20 points)

A source generates

(a) either 10,000 particle pairs in the spin singlet state

$$\frac{1}{\sqrt{2}}(|z\text{-up}\rangle|z\text{-down}\rangle - |z\text{-down}\rangle|z\text{-up}\rangle) = -\frac{1}{\sqrt{2}}(|x\text{-up}\rangle|x\text{-down}\rangle - |x\text{-down}\rangle|x\text{-up}\rangle)$$

(b) or, in a random order, 5,000 pairs in $|z\text{-up}\rangle|z\text{-down}\rangle$ and 5,000 in $|z\text{-down}\rangle|z\text{-up}\rangle$

(c) or, in a random order, 5,000 pairs in $|x\text{-up}\rangle|x\text{-down}\rangle$ and 5,000 in $|x\text{-down}\rangle|x\text{-up}\rangle$.

Alice and Bob are far from each other, and each receives one particle of every pair. By carrying out (local) Stern-Gerlach experiments on their particles and comparing their results afterwards, how can they decide whether the source was of type (a), (b), or (c)?

Exercise 42: Allcock's paradox¹ (30 points)

Allcock considered a "soft detector," i.e., one for which the particle may fly through the detector volume for a while before being detected. An as effective description, Allcock proposed an imaginary potential. For example, for a single particle in 1d and a detector in the right half axis, he considered the Schrödinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = -\frac{\hbar^2}{2m} \frac{\partial^2 \psi}{\partial x^2} - iv1_{x \geq 0} \psi \tag{2}$$

with $v > 0$ a constant. The time evolution then is not unitary.

¹G.R. Allcock: The time of arrival in quantum mechanics II. The individual measurement. *Annals of Physics* **53**: 286–310 (1969)

(a) Derive from (2) the continuity equation

$$\frac{\partial \rho}{\partial t} = -\operatorname{div} j - \frac{2v}{\hbar} 1_{x \geq 0} \rho \quad (3)$$

for $\rho = |\psi|^2$ and $j = \frac{\hbar}{m} \operatorname{Im}[\psi^* \partial \psi / \partial x]$. Eq. (3) is the evolution of the probability density of a particle that moves along Bohmian trajectories and disappears spontaneously (stochastically) at rate $2v/\hbar$ whenever it stays in the region $x \geq 0$. $\|\psi_t\|^2$ is a decreasing function of t and represents the probability that the particle has not been detected (and disappeared from the model) yet.

(b) To obtain an effective description of a hard detector (i.e., one that will detect the particle as soon as it reaches the region $x \geq 0$), Allcock assumed that ψ_0 is concentrated in $\{x < 0\}$ and took the limit $v \rightarrow \infty$, but found that the particle never gets detected ($\|\psi_t\|^2 = \text{const.}$)! That is parallel to the quantum Zeno effect.

Prove the following simplified version: In a 2d Hilbert space \mathbb{C}^2 , let $\psi_0 = (1, 0)$ evolve with the (non-self-adjoint) Hamiltonian

$$H_v = \begin{pmatrix} 0 & 1 \\ 1 & -iv \end{pmatrix}. \quad (4)$$

Then for every $t > 0$, $\psi_t = e^{-iH_v t/\hbar} \psi_0 \rightarrow \psi_0$ as $v \rightarrow \infty$.

Hand in: by Tuesday January 16, 2024, 8:15am

Reading assignment due Thursday January 18, 2024:

A. Einstein, B. Podolsky, N. Rosen: Can Quantum-Mechanical Description of Physical Reality Be Considered Complete? *Physical Review* **47**: 777–780 (1935)